

Liver Transplant Patient Handbook

UCSF Health

UCSF Health

Liver Transplant Phone Number

1 (800) LIVER TX
(548-3789)

Pediatric Phone Number

1 (415) 476-5892

Leave your name, date of birth,
phone number, and a brief message.

Your message will be returned within
one business day.

Table of Contents

Frequently Asked Questions	1	Diuretics	17
Introduction		Anti-ulcer Medications	18
Liver Transplant Service Phone Numbers.	2	High Blood Pressure Medications (Anti-hypertensive)	18
Consulting Other Specialists at UCSF Medical Center	3	Ultrasound	19
UCSF Adult Clinic Phone Numbers	4	CT Scan	19
UCSF Pediatric Clinic Phone Numbers	5	ERCP/PTC	19
Now That You Have Your Liver Transplant	6	Liver Biopsy.	19
Your Local Health Care Provider, the Transplant Team and Your Insurance	6		
Medications and Diagnostic Tests		When You Go Home	
General Information	7	Outpatient Care Guidelines	20
Self-Medication Schedule	7	Liver Transplant Clinic	20
Your Medicine Schedule	7	Adult Clinic	20
Medicine Times.	8	Pediatric Clinic	21
Warnings About Medications.	8	Outpatient Liver Biopsy	21
Immunosuppressive Medications.	9	Daily Routine	23
Tacrolimus (Prograf®).	9	Routine Follow-up Care.	23
Cyclosporine (Neoral®, Gengraf®)	11	Your Local Health Care Provider's Role	24
Mycophenolate Mofetil (CellCept®, Myfortic®)	12	Laboratory Tests and Lab Letters	24
Everolimus (Zortess®)	13	Laboratory Records and Vital Signs Flow Sheets	25
Sirolimus (Rapamune®)	14	Medication Levels	32
Prednisone	15		
Other Medications You May Take After Your Transplant	16	Immunization and Immune Protection	
Antifungal Medications	16	Tuberculosis (TB) testing using purified protein derivative (PPD)	33
Antivirals	16	Immunizations	33
Antibiotics.	17	Pneumonia	33
		Flu vaccine	33

Diphtheria/tetanus toxoid/pertussis (DPT)	33
Tetanus	33
Human Papillomavirus (HPV)	34
Hemophilus B (HIB)	34
Hepatitis A (Hep A)	34
Hepatitis B (Hep B)	34
Inactive Polio Virus Vaccine (IPV)	34
Varicella Zoster (Shingles)	34
Varicella (Chicken Pox)	34
Measles, Mumps, Rubella (MMR)	34
Oral Polio Virus Vaccine (OPV)	34
Immunizations for Pediatric Patients	35

Nutrition Following Liver Transplantation

Short Term Nutrition Management	36
Sodium (salt)	38
Potassium	39
Nutrition for Pediatric Patients Following Liver Transplant	40
Long Term Nutritional Management	40
Low-Fat Cooking Tips	41
Food Safety After Transplant	42

Fitness For Life

How to Start an Exercise Program	43
U .S . Transplant Games	45

Health Maintenance

Skin Cancer	47
Routine Self Exam	47

Acne	47
Dry Skin	48
Cuts and Scratches	48
Increased Hair Growth	48
Hair Care	48
Dental Care	49
Clearance for Dental Care	50
Smoking	51
Alcohol / Illegal Drugs	51
Sexuality	51
Family Planning	51
Return to Work / School/ Disability	52
Travel	53
Other Health Care Concerns	54
Illness and Infection	54
Colds and Flu	54
Herpes Zoster (Shingles)	55
Persistent Headaches	55
Skin Eruptions	55
Thrush	55
Chicken Pox, Measles, and Mumps	55
Emergency Medical Identification	55
Guidelines for Visiting Transplant Children	56

Resources

Getting Back into Life!	58
Adult Support Group	59
Caregiver Support Group	59
Pamphlets / Organizations / Websites	60

Frequently Asked Questions

Medication questions:

- What do I do if I forget to take my anti-rejection (immunosuppressive) medication? **See page 9 under "missed dose" listed under each medication description.**
- What do I do if I need a prescription refill? **Call your pharmacy who will contact the Transplant Office for refill authorization.**
- What do I do if my pharmacy hasn't received prior authorization for a medication? **Call 800-548-3789 and leave a non-urgent message.**
- What over the counter medications can I take? **See page 55.**
- What should I do if am prescribed a medication by another provider? **See page 55.**
- Can I take Tylenol/ acetaminophen, and nonsteroidal anti-inflammatory drugs (NSAIDs)? NSAIDs: ibuprofen, Aleve, Advil, Motrin, Naprosyn, etc. **See page 8.**

Vaccination information:

- Which vaccines can I receive and which should I avoid? **See page 33.**
- What if I am traveling out of the country? **See page 53.**

Labs:

- What do I do when my standing order expires? **Call (800) 548-3789 and leave a non-urgent message. Kaiser insured patients call 888-551-2740 and leave a message for your Kaiser coordinator.**
- How often will I need to have labs drawn? **Frequently right after transplant, then once per month for a lifetime.**
- How can I get my lab results? **Ask your lab how you may also receive lab results. Some labs/ providers have ways you can access your labs on-line.**

- Do my labs need to be drawn while fasting? **No, but you must draw labs first thing in the morning before taking your anti-rejection medications.**

Low Potassium diet:

- What foods should I limit or avoid to keep my potassium level in range? **See page 39.**

Illness and infection:

- What over the counter medications can I take? **See page 54 and page 8.**
- When and who should I call when I am not feeling well? **See page 54.**

Dentist:

- Do I need antibiotics before going to the dentist? **See page 49.**

Medical records

- How do I receive a copy of my medical records? **Call medical records 415-353-2885. The transplant office is not authorized to send medical records.**

Disability and FMLA forms and work release forms: (See page 52)

- How do I get my forms filled out and signed for medical leave? **Call 800-548-3789 and leave a message for your transplant provider.**
- When can I return to work? **See page 52.**

Introduction

Liver Transplant Service Phone Numbers

The Liver Transplant Office is open from 8am to 5pm, Monday through Friday. The Liver Transplant answering service will accept emergency phone calls before and after office hours, on weekends, and all holidays.

Urgent Calls Only

The 24 hour **urgent** transplant number is **415-353-1888**. Pediatric patients should call 415-476-5892. This number is answered 24 hours a day, 7 days a week. If you experience an urgent health concern during non-business hours, please use the urgent transplant phone number. If you feel your concern can wait until the next business day, please call during business hours and ask that your transplant provider (NP/PA) be paged.

Leave the following information when you call the 24 hour urgent transplant number:

- Your name (spell it for the service) and date of birth
- Your phone number(s), including area code
- Brief reason for call

National Emergency Number Is 911

- Call 911 or have a family member or friend take you to the nearest emergency room if you have any of the following:
 - Chest pain
 - Bleeding
 - Trouble breathing
 - Unconsciousness

Non-Emergency Calls

- Reporting lab values is not an emergency.
- Prescription refills are not an emergency.
- Call **1-800-LIVER TX** (548-3789) with questions, or messages for the transplant team. Messages may be left 24 hours a day. Your message will be returned within one business day.
- Requests for routine prescription refills should be called in by your pharmacy to the above number.
- Pediatric patients should call **415-476-5892** or **415-476-8636**.

Consulting Other Specialists

You may be sent to specialists in other departments at the UCSF Medical Center and clinic, so you need to keep a record of the provider's name in each department in order to make an appointment. Keep a list of these providers and their departments for future reference.

Your primary care provider may refer you to specialists within your insurance plan. Include these providers on your list for future reference as well.

Providers in Other Departments

- Cardiologist (Heart)
- Dentist (Teeth)
- Dermatologist (Skin)
- Endocrinologist (Diabetes)
- ENT (Ear/Nose/Throat)
- Ophthalmologist (Eye)
- Gynecologist (Women's health)
- Orthopedist (Bone)
- Primary Care/Internal medicine
- Other

All UCSF Adult Clinic Phone Numbers are in the (415) Area Code

Dental	476-1891
Dermatology	353-7800
Eye	353-2800
ENT	353-2757
Gastroenterology	353-2318
Gynecology/Obstetrics	353-2223
Adult Post Liver Transplant Clinic appt	353-9130
Pediatric Liver Transplant Clinic appt.	353-2813
Medicine	353-4624
Neurology	353-2273
Orthopedic	353-2808
UCSF Pharmacy	353-1544
Urology	353-2555

Other Services:

Social Work (message line)	800-548-3789
--------------------------------------	--------------

Pediatric Clinics at UCSF Mission Bay

Phone Numbers All are in the (415) Area Code

Cardiology	353-2008
Dental.	476-1891
Dermatology.	353-7800
Eye (ophthalmology)	353-2800
ENT (otolaryngology).	353-2757
Gastroenterology/Liver	353-7337
Gynecology/Obstetrics	353-9600
Neurology.	353-2273
Orthopedic	353-2808
Renal (kidney).	353-7337
Urology.	353-2200
Surgical Specialties.	353-4287

Other Services:

Social Work Office	353-2655
Pediatric Infusion Center.	353-2584

Now That You Have Your Liver Transplant

This booklet was prepared to help you learn what you need to know about your new liver. Although we are here to help you, it is **your responsibility** to learn how to take care of yourself and your new liver. You are the most important person in your health care team so you need to know your medications, lab values, and the routine follow-up care needed.

In the hospital, your nurses will begin to teach you promptly. At first it may seem like a lot of information, but it will get easier. It's normal to have questions after you are discharged from the hospital. This handbook is a quick reference guide and a "starting point" for answering your questions. Always ask your nurse or the transplant team any questions you might have. We are here to help you.

Write your questions down so you do not forget them. There is space to write your questions at the end of this handbook.

Your Local Health Care Provider, The Transplant Team and Your Insurance

Your local health care providers and the transplant team are committed to providing care for you in the best way possible. This commitment includes providing care within the guidelines outlined by your insurance company. After the first hospital discharge and follow-up appointment, you may need to get pre-authorization from your primary care provider for clinic visits or procedures with the transplant team.

It is your responsibility to get any necessary authorizations that have been requested and obtain these from your primary care physician or you may be billed for some charges. You may be able to have some testing such as labs and liver biopsies done and imaging studies (CTs/CAT, etc.) are done in your local area. Know as much as you can about the pre-authorization requirements of your insurance company, as well as any special features about your coverage.

Examples: When do you need a pre-authorization?

- *Which pharmacies are "preferred" within your insurance system?*
- *Which labs are "preferred" within your insurance system?*

Medications and Diagnostic Tests

General Information

You will take several medications after your liver transplant for the rest of your life. Each medication has been individually prescribed for you, and it is very important that you take your medications as you are instructed. We want you to be responsible for taking your own medications. We also want children to be involved in taking their own medications. There is more information about these medications later in this chapter.

Self-Medication Schedule

After your transplant, your nurse will set up a medication schedule for you. This medication schedule is set up to help you learn about the medications you are taking, when to take them, what they look like, how much to take, what they are for, and their side effects. When you begin to take your own medicine after surgery, your nurse must always check your pills before you take them. When you go home from the hospital you may be taking 10–14 different medications. These include medications you will take for the rest of your life and other medications that will stop in weeks to months.

Your Medicine Schedule

You will be given a copy of your medicine schedule with a list of all your medications to keep at your bedside. Record any changes on this schedule in pencil.

- Never follow the directions on the bottle as these directions may change.
- Write any dosage changes, medication changes, or new medicines on your schedule.
- Remember to bring your medicine schedule when you come to the clinic or to the hospital. Also bring the medication bottles if you have any questions; for pediatric patients, always bring medication bottles in addition to your schedule.
- Never take any new medications without letting the UCSF Liver Transplant Office know. Many medications change the effects of your immunosuppressive medications. These include but are not limited to: Erythromycin (Ery-tabPediazole®), Azithromycin (Zithromax®), Clarithromycin (Biaxin®), Voriconazole, Fluconazole, Posaconazole (Noxafil®), Ibuprofen (Advil®, Motrin®), Naproxen (Aleve®), Phenytoin (Dilantin®), Phenobarbital, and herbal supplements such as St. John's Wort or Echinacea .

Medicine Times

You must take your pills at the same time every day. In the hospital, we use standard times of 9am, 12 noon, 6pm, and bedtime. When you go home, you should use times that fit your lifestyle but the medications should be taken at the same time each day. Your Tacrolimus or Cyclosporine should be taken 12 hours apart from each other, for example 9 AM and 9 PM. Your nurses will help you set up a system.

Warnings About Medications

Never:

- Never change or stop any immunosuppressive medication the transplant team has prescribed without checking with us first.
- You must never forget to take your medications.
- Never take Motrin, Ibuprofen, Advil, Naproxen, Aleve, and other Non-Steroidal Anti-inflammatory drugs (also called NSAIDS), as the combined effect with Tacrolimus (Prograf) or Cyclosporine (Neoral, Gengraf) can cause kidney damage. Drugs included in this category may be found in over the counter common cold remedies. Please check labeling to ensure that you use a product that does not contain these medications.

Always:

- Use acetaminophen (example: Tylenol®) only for headache or pain (maximum dose 2000 mg/day). Parents should be sure the dose is correct for their child's weight. Remember, Tylenol is also in some narcotics, for example Vicodin and Norco. The total maximum dose of 2000 mg/day includes prescription and over the counter medication.
- Check with your pharmacist or transplant team if you have any questions about any of your medications.
- Store your medicine out of the reach of small children. Remember that your medicine could be deadly if accidentally taken by a child. Pharmacies put all prescriptions in child proof containers. If you do not want child proof containers, ask your pharmacist for regular containers.
- Plan ahead: always call for prescription refills at least 1–2 weeks before you run out of medication.
- Call **1-800-LIVER TX** (1-800 548-3789) with any questions you have about medications. We make all attempts to get back to you within 24 hours. Office hours: Monday–Friday, 8 AM–5 PM .

Immunosuppressive Medications

Because your new liver is “foreign” to your body, your immune system will try to reject it. Therefore, it is necessary to suppress your immune system to protect your new liver. The drugs described in this chapter prevent and/or treat rejection of your new liver. Immediately after your transplant, when you are most likely to reject your new liver, you will be taking very high doses of these medications. As the risk of rejection decreases, the doses will be reduced. When the doses are high, you may experience some side effects.

Remember that not every patient experiences all of these side effects, but they occur often enough that you should know about them. Do not hesitate to tell the transplant team about any side effects or problems you may have. We can often help to lessen the side effects.

Tacrolimus (Prograf®), Cyclosporine (Neoral®, Gengraf®), Mycophenolate Mofetil (CellCept®), Mycophenolic acid (Myfortic®), Everolimus (Zortress®), Sirolimus (Rapamune®), Prednisone

These are the immunosuppressive drugs prescribed to prevent rejection of your transplanted liver. You will be on a combination of these drugs at different times after your transplant. The drug or the dose of each of these drugs may change depending on your particular condition. Your transplant team will help you with changes in your medications while you are in the hospital. After you leave the hospital, you will receive phone calls to adjust the dosages based on your lab results. Write these changes on your medication schedule immediately along with the date of the change.

<i>Tacrolimus</i> Prograf®	
Purpose	<ul style="list-style-type: none">▪ Tacrolimus is an immunosuppressant medication that helps prevent or treat rejection.
Dose	<ul style="list-style-type: none">▪ The tablets are available in 0.5 mg, 1 mg, and 5 mg strengths; liquid is 0.5 mg/ml. The liquid must be specially made by your pharmacist. The dose is adjusted by your laboratory values. It is very important to have your labs drawn 12 hours after your last Tacrolimus dose and before you take your morning medications for proper adjustment of this medication.
When to Take	<ul style="list-style-type: none">▪ AM Dose: take with your breakfast, after your lab work is completed (usually around 9am).▪ PM Dose: 12 hours after your AM dose (usually around 9pm).

Tacrolimus

Prograf®

Missed Dose	<ul style="list-style-type: none">▪ If you miss a dose, make it up as soon as you remember. But don't take two doses at one time without contacting the transplant office.
Precautions	<ul style="list-style-type: none">▪ Tacrolimus (Prograf) can be altered by some medications. Before you take any new drugs including over the counter medications and medications prescribed by your local provider, please contact the transplant office. It is recommended that mothers do not breast feed when taking this medication. You should avoid grapefruit, pomegranate, pomelo, starfruit, and all juices containing these fruits since they can interfere with your Tacrolimus level.
Common Side Effects	<p><i>The side effects of Tacrolimus are similar to those of Cyclosporine.</i></p> <ul style="list-style-type: none">▪ Headaches▪ Diarrhea▪ Seizures▪ High blood pressure▪ Tremors▪ Rash▪ Mental status changes▪ Decreased magnesium▪ Insomnia▪ High blood sugar▪ Anemia▪ High potassium▪ Nausea/vomiting▪ Decreased kidney function▪ Decreased appetite▪ Leg cramps▪ Hair Loss

Cyclosporine (modified)

(Neoral[®] or Gengraf[®]):

<p>Purpose</p>	<ul style="list-style-type: none"> ▪ Cyclosporine is an immunosuppressant medication that helps prevent rejection.
<p>Dose</p>	<ul style="list-style-type: none"> ▪ Cyclosporine capsules are available in 25 mg and 100 mg strengths; liquid is 100 mg/ml. Do not combine the brands of cyclosporine you take. In other words, prescriptions for both 25 and 100 mg capsules should be consistently filled with the same brand of Cyclosporine modified. Call the transplant team with any questions. The dose is adjusted by your laboratory values. It is very important to have your labs drawn 12 hours after your last Cyclosporine dose and before you take your morning medications for proper adjustment of this medication.
<p>When to Take</p>	<ul style="list-style-type: none"> ▪ AM Dose: take with your breakfast, after your lab work is completed (usually around 9am). ▪ PM Dose: 12 hours after your AM dose (Usually around 9pm).
<p>Missed Dose</p>	<ul style="list-style-type: none"> ▪ If you miss a dose, make it up as soon as you remember. But don't take two doses at one time without contacting the transplant office.
<p>Precautions</p>	<ul style="list-style-type: none"> ▪ Cyclosporine can be altered by some medications. Before you take any new drugs including over the counter medications and medications prescribed by your local provider, please contact the transplant office. It is recommended that mothers do not breast feed when taking this medication. You should avoid grapefruit, pomegranate, pomelo, starfruit, and all juices containing these fruits since they can interfere with your Cyclosporine level.
<p>Common Side Effects</p>	<ul style="list-style-type: none"> ▪ Headaches ▪ Loss of Appetite ▪ Tremors ▪ Diarrhea ▪ Seizures ▪ Nausea/vomiting ▪ Nightmares ▪ Rash ▪ Memory loss ▪ Hair growth ▪ Fluid retention ▪ Gum swelling ▪ Decreased kidney function ▪ Increased risk of infection ▪ Elevated potassium levels ▪ Joint discomfort ▪ High blood pressure ▪ Leg cramps ▪ Hot flashes

Mycophenolate

Cellcept® or Myfortic®

Purpose	<ul style="list-style-type: none">▪ Mycophenolate is an immunosuppressant medication that helps prevent rejection. Myfortic is the enteric coated form of Mycophenolate that can have less stomach and bowel side effects.
Dose	<ul style="list-style-type: none">▪ Mycophenolate is available in 250 mg strength and 500 mg tablets. Note that tablets are also available in 500 mg strength, but we will write your prescriptions using 250 mg strength so it will be easier to make dosage changes. Liquid comes in 200 mg/ ml strength. Be sure to check the bottle you get from your pharmacy to make sure you have the right dose. Myfortic is available in 180 mg and 360 mg tablets. Do not crush or chew Myfortic tablets. <p><i>Note: the doses of Mycophenolate and Myfortic are not the same due to the way they are absorbed in the body.</i></p>
When to Take	<ul style="list-style-type: none">▪ AM Dose: Take with your breakfast (usually around 9am).▪ PM Dose: 12 hours after your AM dose (usually around 9pm).
Missed Dose	<ul style="list-style-type: none">▪ If you miss a dose, make it up as soon as you remember. But don't take two doses at one time without contacting the transplant office.
Precautions	<ul style="list-style-type: none">▪ Antacids such as Maalox® and Mylanta® decrease the absorption of Mycophenolate. You should not take these while on Mycophenolate. We do recommend that you continue to take a medication such as Omeprazole (Prilosec®), Pantoprazole (Protonix®), Lansoprazole (Prevacid®), etc . while you are on Mycophenolate. You should not take Mycophenolate (Cellcept® or Myfortic®) for at least 3 months before conceiving children, please check with the transplant office if you are considering having children.▪ Please check with the transplant office before you consider becoming pregnant.▪ It is recommended that mothers do not breast feed when taking this medication.
Common Side Effects	<ul style="list-style-type: none">▪ decreased white blood count▪ diarrhea▪ abdominal pain▪ ulcers▪ inflammation of the pancreas

Everolimus

Zortress®

Purpose	<ul style="list-style-type: none">▪ Everolimus is an immunosuppressant medication that helps prevent or treat rejection.
Dose	<ul style="list-style-type: none">▪ The tablets are available in 0.25 mg, 0.5 mg, and 0.75 mg strength. The dose is adjusted to your lab results. It is important to have labs drawn 12 hours after your last dose prior to your next dose.
When to Take	<ul style="list-style-type: none">▪ AM Dose: Take with your breakfast (usually around 9am).▪ PM Dose: 12 hours after your AM dose (usually around 9pm).
Missed Dose	<ul style="list-style-type: none">▪ If you miss a dose, make it up as soon as you remember. But don't take two doses at one time without contacting the transplant office.
Precautions	<ul style="list-style-type: none">▪ Please check with the transplant office before you consider becoming pregnant.▪ It is recommended that mothers do not breast feed when taking this medication.▪ You should avoid grapefruit, pomegranate, pomelo, starfruit, and all juices containing these fruits since they can interfere with your everolimus level.
Common Side Effects	<p>The most common side effects of Everolimus are changes in blood counts: anemia (low red blood counts), low platelets, and low white blood cells. Other side effects may include:</p> <ul style="list-style-type: none">▪ Mouth sores▪ Rash▪ Poor wound healing▪ Headaches▪ Diarrhea▪ Tremors▪ Swelling▪ Kidney damage <p><i>*Alert: you will need to be taken off of Everolimus approximately one month prior to any surgery due to delayed wound healing. Please inform your transplant team of any upcoming surgery.</i></p>

Sirolimus

Rapamune®

Purpose	<ul style="list-style-type: none">▪ Sirolimus is an immunosuppressant medication that helps prevent or treat rejection.
Dose	<ul style="list-style-type: none">▪ The tablets are available in 0.5 mg, 1 mg, and 2 mg strength. This is a once per day dosing schedule. The dose is adjusted to your lab results. It is important to have labs drawn 24 hours after your last dose prior to your next dose. Liquid is available as 1 mg/mL and should be mixed with orange juice or water only in a glass or plastic container.
When to Take	<ul style="list-style-type: none">▪ If you also take Cyclosporine (Neoral® or Gengraf®) as another immunosuppressant medication, you must take the Sirolimus 4 hours after the Cyclosporine dose as these drugs should NOT be taken at the same time.▪ If you take Tacrolimus (Prograf), it is ok to take Sirolimus at the same time.
Missed Dose	<ul style="list-style-type: none">▪ If you miss a dose, make it up as soon as you remember. But don't take two doses at one time without contacting the transplant office.
Precautions	<ul style="list-style-type: none">▪ Please check with the transplant office before you consider becoming pregnant.▪ It is recommended that mothers do not breast feed when taking this medication.
Common Side Effects	<p>The most common side effects of Sirolimus are changes in blood counts: anemia (low red blood counts), low platelets, and low white blood cells. Other side effects may include:</p> <ul style="list-style-type: none">▪ Mouth sores▪ Rash▪ Poor wound healing▪ Headaches▪ Diarrhea▪ Tremors▪ Swelling▪ Kidney damage <p><i>*Alert: you will need to be taken off of Sirolimus approximately one month prior to any surgery due to delayed wound healing. Please inform your transplant team of any upcoming surgery.</i></p>

Prednisone

Purpose	<ul style="list-style-type: none"> This is an immunosuppressant medication that is used to prevent or treat rejection.
Dose	<ul style="list-style-type: none"> Adults: Use only 5 mg and 10 mg tablets unless first contacting the transplant office. Children: Prednisone solution is available in the following concentrations: 1 mg/ml; 5 mg/ml; and 15 mg/5 ml. Be sure you read the label. If you have questions, call the transplant office. Also, Prednisone is available in 1 mg and 5 mg tablets for older children who are able to swallow pills.
When to Take	<ul style="list-style-type: none"> Prednisone is taken once per day in the morning. You will be given a schedule to follow that tells you how much prednisone to take every day. You will be on a higher dose right after transplant and will decrease the daily amount by following the medication schedule. By the time you are two months after transplant, you will usually be on 5 mg once a day taken in the morning. Many children will stop taking Prednisone by the time they are 2 to 3 months after transplant. Take with food to avoid upset stomach. The dose for children varies depending on their weight.
Missed Dose	<ul style="list-style-type: none"> If you miss a dose, make it up as soon as you remember. But, don't take two doses at one time without contacting the transplant office.
Precautions	<ul style="list-style-type: none"> It is important that you never stop taking your Prednisone abruptly. This could cause severe side effects. If you are a diabetic it is important to remember that as your Prednisone dose decreases, your insulin requirements will also decrease.
Common Side Effects	<ul style="list-style-type: none"> Mood changes or mood swings Depression Euphoria Irritability Vision changes Thrush Acne Increased risk of infection Stomach upset High blood sugar Moon face Cataracts Sensitivity to the sun High blood pressure Joint discomfort Nausea/vomiting Muscle wasting Ulcers Increased appetite

Other Medications You May Take After Your Transplant

Anti Fungal Medications

Fluconazole (Diflucan®)

Purpose	<ul style="list-style-type: none"> Prevents or treats fungal (Candida) infections often found in the mouth.
Dose	<ul style="list-style-type: none"> This medication is taken one time weekly. The dose is generally 100 mg. This medication will be discontinued once your prednisone dose reaches 5 mg daily, usually two months after your transplant.
When to Take	<ul style="list-style-type: none"> Once per week.
Missed Dose	<ul style="list-style-type: none"> If you miss a dose, make it up as soon as you remember. But, don't take two doses at one time.
Precautions	<ul style="list-style-type: none"> Never take this medication more than once per week unless under specific instruction with the transplant department due to the interaction with your transplant medications.

Antivirals

Valgancyclovir (Valcyte®) and Acyclovir (Zovirax®).

Purpose	<ul style="list-style-type: none"> Prevents or treats viral infections such as CMV, cold sores, herpes zoster, and chicken pox.
Dose	<ul style="list-style-type: none"> Valgancyclovir is available in 450 mg tablets or in liquid form (50 mg/ml). The adult dose is usually 900 mg per day for the first 3 to 6 months after transplant depending on your kidney function. Acyclovir is available in 200 mg capsules, 800 mg tablets or in liquid form (200 mg/5 ml). The adult dose is usually 800 mg 4 times a day for 3 months after transplant. The children's dose is based on body weight. Adults are usually treated for 3 to 6 months after transplant. Children are usually treated for 6 to 12 months after transplant.
Missed Dose	<ul style="list-style-type: none"> Take as soon as you remember. Never double dose.
Side Effects	<ul style="list-style-type: none"> May include decreased white count. Dosage may be decreased for people with kidney problems.

Antibiotics

Trimethoprim/sulfamethoxazole (Bactrim® or Septra®), dapsone, pentamidine

Purpose	<ul style="list-style-type: none">▪ Treats or prevents bacterial infections, such a pneumocystis pneumonia.
Dose	<ul style="list-style-type: none">▪ For Trimethoprim/Sulfamethoxazole (Bactrim® or Septra®) the adult dose is 1 tablet every Monday, Wednesday, Friday; Dapsone is 50–100 mg once a day; Pentamidine is a once a month inhalation treatment. Pediatric doses are based on body weight.
Missed Dose	<ul style="list-style-type: none">▪ Take as soon as you remember. Never double dose.
Side Effects	<ul style="list-style-type: none">▪ Side effects of trimethoprim/sulfamethoxazole include a rash, decreased white count, nausea, vomiting, diarrhea, sun sensitivity.▪ Side effects of Dapsone include decreased red blood cell count, fatigue, and shortness of breath.▪ Side effects of pentamidine include coughing.▪ You will be on this medication for about 1 year after transplant.
Precautions	<ul style="list-style-type: none">▪ Some patients may not be able to take Septra/Bactrim® (sulfa medications). For those patients, Dapsone will be prescribed. Another alternative is Pentamidine which is given monthly as a respiratory inhalation treatment.

Diuretics – “water pills”

Furosemide (Lasix®)

Purpose	<ul style="list-style-type: none">▪ To remove excess fluid from your body and sometimes for blood pressure management.
Dose	<ul style="list-style-type: none">▪ Varies according to your weight, kidney function, potassium level, urine output, and blood pressure. Try to take doses in the morning or early afternoon to avoid frequent urination in the middle of the night.
Missed Dose	<ul style="list-style-type: none">▪ Take as soon as you remember. Never double dose.
Side Effects	<ul style="list-style-type: none">▪ Weakness, muscle cramps, light-headedness, dizziness.

Anti Ulcer Medications

Omeprazole (Prilosec[®]), Esomeprazole (Nexium[®]), Pantoprazole (Protonix[®]), Lansoprazole (Prevacid[®])

Purpose	<ul style="list-style-type: none">▪ To prevent ulcers. Ulcers can develop with high doses of Prednisone and while taking Mycophenolate (Cellcept[®] or Myfortic[®]).
Dose	<ul style="list-style-type: none">▪ Dose and type of medication is based on individual need and body weight.
Missed Dose	<ul style="list-style-type: none">▪ Take as soon as you remember. Never double dose.
Side Effects	<ul style="list-style-type: none">▪ Side effects are uncommon.

High Blood Pressure Medications (Anti-hypertensives)

Nifedipine (ProcardiaXL[®]), Metoprolol, Amlodipine (Norvasc[®])

Purpose	<ul style="list-style-type: none">▪ To control your blood pressure.
Dose	<ul style="list-style-type: none">▪ Dose and type of medication is based on individual need and body weight.
Missed Dose	<ul style="list-style-type: none">▪ Take as soon as you remember. Never double dose.
Side Effects	<ul style="list-style-type: none">▪ Dizziness, nausea, headache, flushing, changes in heart rate.

Diagnostic Tests

The list below describes the purposes of a variety of diagnostic procedures you may undergo after you have had your liver transplant.

Ultrasound

This is sometimes done the day after transplant to make sure the main blood vessels leading to the liver are functioning normally. An ultrasound can also be used to check for abnormal fluid collection (blood, bile) in the abdomen and to localize position for biopsies in children. A gel is applied on your abdomen and a transducer (a wand-like instrument) is passed over the abdomen. Images of your liver will appear on a monitor.

CT Scan

A type of x-ray that looks at your liver from several angles to check for infection or fluid collections.

After you drink a liquid dye, pictures are taken while you lay still. Children are usually given something to relax them before this procedure is performed.

ERCP/PTC

There are different types of x-rays that check bile ducts for leaks, blockages, or other potential problems. ERCP (endoscopic retrograde cholangiogram) is done by placing a tube down your throat through which a catheter is inserted into a bile duct. PTC (percutaneous transhepatic cholangiogram) is done by inserting a needle into the liver through the abdomen, then placing the catheter in the bile ducts. You will be given medication to relax you before the ERCP or PTC. Children usually have these tests done under anesthesia.

Liver Biopsy

A liver biopsy checks for rejection, hepatitis, and other possible problems. You may need a liver biopsy if your liver numbers from labs go up. The right side of the abdomen is numbed, a biopsy needle is inserted, and a small piece of liver tissue will be removed. You must lay down on your right side for at least 1 hour and stay in bed for 4 hours after this procedure. You may not take food or fluids during this time.

You must have transportation arranged by a friend or relative. Public transportation (cabs, buses) is not allowed. Your procedure will be cancelled if you do not have adequate supervised transportation.

You must be off blood thinners, such as warfarin (Coumadin) or aspirin or Ibuprofen prior to the biopsy. See *page 23 for guidelines*.

Infants and children are given mild sedation during the biopsy and are admitted to the hospital for the procedure.

When You Go Home

Outpatient Care Guidelines

Call the Transplant Emergency phone number, (415) 353-1888, or your local healthcare provider for temperatures over 100.4°F (38°C). The most important thing you can do to ensure your long term health is to recognize and report any changes in the way you are feeling. Please do not try to “self-diagnose.” Because of your antirejection or immunosuppressive medications, symptoms may need urgent treatment. Pediatric patients should call 415-476-5892.

Liver Transplant Clinic

You will be followed by a Transplant Center for the rest of your life. You will be seen on a regular basis in the Liver Transplant clinic. Children will be followed in the Pediatric GI/Transplant clinic.

Please bring your medicine schedule and medications for the entire day with you. Remember to check in at the clinic desk when you arrive. Follow-up appointments may be made at the desk when you leave or you may call 415-353-9130. Pediatric patients can call 415-353-2813.

Labs

You need to have your blood drawn twice weekly the first month after your transplant and once weekly the next month following your transplant. After that you will probably have your labs done twice a month, then once a month. You can have your blood drawn at a local laboratory who will fax your lab results to the Liver Transplant Office. *Fax for adults: 415-353-8917. Fax for children: 415-476-1343.*

Adult Clinic

The adult Liver Transplant Clinic is on the 7th floor of the Ambulatory Care Center (ACC) and meets every Tuesday from 8am–12pm and Thursday afternoon from 1:30pm–3:30pm. Call 415-353-9130 to make appointments at the adult clinic. You will be given a specific time for your appointment. If your insurance requires authorization for clinic visits, it is your responsibility to arrange with your primary care physician and have this approved before the visit.

Outreach Clinic

UCSF has monthly adult outreach clinics in Fresno and Modesto, and a bimonthly outreach clinic in Walnut Creek. To schedule an appointment at any of the UCSF outreach clinics, please call 415-353-2156.

Pediatric Clinic

The Pediatric GI/Liver Clinic is at the Mission Bay Campus/Benioff Children's Hospital on the 6th floor of the Gateway Medical Building and meets Thursdays from 8am–12pm. Call 415-353-2813 to make appointments at the pediatric clinic.

Outpatient Liver Biopsy

The frequency of outpatient liver biopsies will depend on what types of problems you have. Biopsies may be done to evaluate for rejection or hepatitis. Do not eat or drink anything after midnight before the biopsy, except for your medications. You must have a current set of labs drawn one week prior to your biopsy. You will be instructed to stop taking aspirin (also don't take Motrin, Advil or Ibuprofen) 5 days before and 5 days after your biopsy. If you are taking any other blood thinner such as warfarin (Coumadin) your biopsy will require careful coordination. Be sure to take your morning medications the day of the biopsy. If you are diabetic, ask about adjusting your insulin dosage for that morning. You will stay in the surgery center for about 6 hours after the biopsy is performed.

You must have transportation arranged by a friend or relative. Public transportation (cabs, buses) is not allowed. Your procedure will be cancelled if you do not have adequate supervised transportation.

Infants and small children may have liquids up until 4 hours before the biopsy. Infants and small children are always admitted to the hospital for this procedure. Check with your physician or transplant nurse practitioner for questions.

Location

Adult outpatient liver biopsies: These are performed at the PPU (Peri-Procedural Unit), 505 Parnassus, 6th floor. You should report there at 7:30am the morning of your biopsy.

Your local gastroenterologist may perform this procedure at your local hospital and the slides will be sent to our pathology department for review.

Pediatric liver biopsies are generally done at the Benioff Children's Hospital. Most children are admitted in the morning of the procedure and spend the night. If rejection is seen on the biopsy, children may remain hospitalized for a few days to a week, depending on the treatment. An ultrasound is performed to mark the spot for biopsy prior to the procedure. Be sure to check with your transplant provider for arrival time and additional instructions.

After the first few months, your local gastroenterologist may perform this procedure at your local hospital and the slides will be sent to us for review.

Adult Liver Biopsy Info

Where: Please check in to the Admitting Office – 505 Parnassus Avenue, Ground Floor, room M140. Once checked in, you will be directed to the Peri-Procedural Unit. Take the Moffitt elevators up to the 6th floor. Follow the signs for the Peri-Procedural Unit, and check in at the nurses' station on 6 East.

Peri-Procedural Unit phone number: 415-353-9123.

Time: 7:30am check in

Instructions:

- No Food or Drink after Midnight. Little water to take medications.
- No Aspirin or Coumadin or NSAIDS 1 week prior to appointment.
- Do standing order and additional labs 1 week prior to appointment.

Daily Routine

Follow a daily routine similar to the one outlined below after you go home.

Mornings

1. Weigh yourself each morning after going to the bathroom. Record your weight in your handbook.
2. Take and record your temperature.
3. Take and record your blood pressure if you are taking anti-hypertensive medications.
4. Take your morning medications, except as described in number “5” below.
5. Bring your Cyclosporine (Gengraf® or Neoral®) or Tacrolimus (Prograf®) or Sirolimus (Rapamune®) with you to the lab on the days you have your Cyclosporine, Tacrolimus, or Sirolimus level drawn; then take your morning Cyclosporine, Tacrolimus, or Sirolimus dose **after** your blood is drawn.

Noon

6. Take your medications as ordered by the Transplant Team.

Nights (between 7–9pm):

7. Take your bedtime medications.
8. Set out your medications for the next day. Use of a multi-chambered plastic pill box may be helpful.

Routine Follow-up Care

The first 60–90 days after your transplant are the time of highest risk for rejection and infection. You will need to have frequent blood tests. Your vital signs (pulse, blood pressure, temperature) must be checked often during the first three months, and after this time your vital signs can be checked as needed. The risk of rejection usually decreases after the first 60–90 days. The frequency of blood tests and vital signs can also be reduced. Your transplant providers will let you know when you can decrease the frequency of your laboratory tests.

Be sure to keep the Transplant Office informed of your correct address and phone number(s), and any changes in your health care provider’s name, address, or phone number(s).

It is also important to tell us whom to contact in case of an emergency. You will be given a form to complete. Be sure to return the completed form to us.

If you are an outpatient and need urgent medical care or need to see a health care provider at UCSF, call 415-353-1888 (days, evenings, and holidays) or go to your local emergency room.

Pediatric patients should call 415-476-5892.

Do not come to the Patient Care Unit or Clinic unless told to do so by the Transplant Team.

Your Local Health Care Provider's Role

Your local health care provider may be the one who took care of you before you came to UCSF for your liver transplant, or who referred you to us. Your visit to this provider is very important and should occur no later than one month after you return home. You should also continue seeing those specialists who were involved in your care before you received a liver transplant. For example: your pediatrician for children, your cardiologist for heart disease, your gynecologist for women, your ophthalmologist for eye care, and/or your family provider for general health care needs, etc.

After the immediate post transplant period (in other words, more than 6–12 weeks after your release from the hospital), your first contact for illnesses should be with your local provider. After he or she has seen you, your provider should contact the Transplant Office with any questions.

Remember – Contact the Transplant office during business hours if a new medication is prescribed by your local health care provider. Many medications can have significant interactions with your antirejection or immunosuppressive medications. Inform the Transplant Office of any changes in your health care provider's name, address, or phone number(s).

Laboratory Tests and Lab Letters

Blood tests need to be done frequently for about 90 days after your transplant. They are done every day while you are hospitalized and twice weekly after you go home.

Preferred lab locations are Kaiser, Lab Corp or Quest labs as we receive these results promptly.

Infants and small children are often discharged after transplant with a central line for blood draws. The central line is an intravenous catheter placed at the time of transplant in the front of the chest. Your nurse will show you how to take care of it before you go home, and you might have a home health nurse checking in with you after you are discharged from the hospital. When blood draws are decreased to once a week, the central line is typically removed. This requires a “come and go” outpatient surgical procedure.

Regular laboratory days after you go home are Mondays and Thursdays. You will take home a laboratory order that lists all the tests that need to be done and how often. Give this order to your laboratory.

Your lab should fax results to the Transplant Office and to your local provider. If you would like a copy of your laboratory results, please ask your laboratory for these.

Lab results should be faxed to the Transplant Office at 415-353-8917 (adult) or 415-476-1343 (pediatrics).

Laboratory Records and Vital Signs

Date:

Hematocrit (Hct)							
Hemoglobin (Hgb)							
WBC							
Platelets (Plts)							
BUN							
Creatinine (Cr)							
Sodium (Na)							
Potassium (K)							
Glucose							
Bili - Total							
SGOT/AST							
ALK P'tase							
SGPT/ALT							
Cyclosporine or Prograf level							
Sirolimus or Everolimus level							
Weight							
Temp							
Blood Pressure							

Laboratory Records and Vital Signs

Date:

Hematocrit (Hct)							
Hemoglobin (Hgb)							
WBC							
Platelets (Plts)							
BUN							
Creatinine (Cr)							
Sodium (Na)							
Potassium (K)							
Glucose							
Bili - Total							
SGOT/AST							
ALK P'tase							
SGPT/ALT							
Cyclosporine or Prograf level							
Sirolimus or Everolimus level							
Weight							
Temp							
Blood Pressure							

Laboratory Records and Vital Signs

Date:

Hematocrit (Hct)							
Hemoglobin (Hgb)							
WBC							
Platelets (Plts)							
BUN							
Creatinine (Cr)							
Sodium (Na)							
Potassium (K)							
Glucose							
Bili - Total							
SGOT/AST							
ALK P'tase							
SGPT/ALT							
Cyclosporine or Prograf level							
Sirolimus or Everolimus level							
Weight							
Temp							
Blood Pressure							

Laboratory Records and Vital Signs

Date:

Hematocrit (Hct)							
Hemoglobin (Hgb)							
WBC							
Platelets (Plts)							
BUN							
Creatinine (Cr)							
Sodium (Na)							
Potassium (K)							
Glucose							
Bili - Total							
SGOT/AST							
ALK P'tase							
SGPT/ALT							
Cyclosporine or Prograf level							
Sirolimus or Everolimus level							
Weight							
Temp							
Blood Pressure							

Laboratory Records and Vital Signs

Date:

Hematocrit (Hct)							
Hemoglobin (Hgb)							
WBC							
Platelets (Plts)							
BUN							
Creatinine (Cr)							
Sodium (Na)							
Potassium (K)							
Glucose							
Bili - Total							
SGOT/AST							
ALK P'tase							
SGPT/ALT							
Cyclosporine or Prograf level							
Sirolimus or Everolimus level							
Weight							
Temp							
Blood Pressure							

Laboratory Records and Vital Signs

Date:

Hematocrit (Hct)							
Hemoglobin (Hgb)							
WBC							
Platelets (Plts)							
BUN							
Creatinine (Cr)							
Sodium (Na)							
Potassium (K)							
Glucose							
Bili - Total							
SGOT/AST							
ALK P'tase							
SGPT/ALT							
Cyclosporine or Prograf level							
Sirolimus or Everolimus level							
Weight							
Temp							
Blood Pressure							

Laboratory Records and Vital Signs

Date:

Hematocrit (Hct)							
Hemoglobin (Hgb)							
WBC							
Platelets (Plts)							
BUN							
Creatinine (Cr)							
Sodium (Na)							
Potassium (K)							
Glucose							
Bili - Total							
SGOT/AST							
ALK P'tase							
SGPT/ALT							
Cyclosporine or Prograf level							
Sirolimus or Everolimus level							
Weight							
Temp							
Blood Pressure							

Medication Levels

If you are taking Tacrolimus (Prograf®) or Cyclosporine (Neoral®, Gengraf®, Sandimmune), Everolimus (Zortress®) or Sirolimus (Rapamune®), your blood must be tested frequently to measure the drug level in your blood. Do not take your morning dose of medication before your labs. Bring your Tacrolimus, Cyclosporine, Everolimus, or Sirolimus with you to the lab and take your morning dose after your blood is drawn.

Immunization and Immune Protection

It is extremely important for pediatric liver transplant patients and children of transplant recipients to receive appropriate immunizations to prevent disease. However, after transplant the recipient's immune system is compromised due to the immunosuppressive medications that must be taken to prevent rejection.

Therefore, liver transplant patients **should not receive any live virus vaccines**, such as measles, mumps, rubella (MMR), small pox, chicken pox, shingles (Zostavax[®]), nasal flu mist, and oral polio vaccine (OPV).

Transplant patients can receive other vaccines such as diphtheria, tetanus toxoid, pertussis (DPT), and inactive polio virus vaccine (IPV), or human papilloma virus (HPV).

The following highlights important information about immunizations and the liver transplant patient and family.

Tuberculosis (TB) testing using purified protein derivative (PPD)

TB testing using PPD may be performed on liver transplant patients and household members.

Immunizations

If you are planning a trip to a foreign country that requires any immunizations, you may need to have your health care provider write a letter to the Passport Bureau stating that you cannot receive certain vaccines. If you have questions about vaccinations or precautions to take, ask your primary care provider. You may be referred to a travel clinic. Another resource you may contact is the Overseas Medical Center, phone 415-982-8380.

Inactivated Vaccines: (These are OK to take)

Pneumonia Vaccine

Flu Vaccine

Inactive virus: Transplant patients and household members may receive vaccine.

Never receive the nasal flu mist—this is a live vaccine.

Diphtheria/tetanus toxoid/pertussis vaccine (DPT or Tdap)

Inactive virus: Transplant patients and household members may receive vaccine.

Tetanus

Toxoid: Generally given every 10 years. Liver transplant patients and household members may receive tetanus shots.

Human Papillomavirus (HPV)

Inactive Virus: Transplant patients and household members may receive vaccine.

Hemophilus B (HIB)

Polysaccharide: Transplant patients and household members may receive HIB.

Hepatitis A (Hep A)

Inactive virus: Transplant patients and household members may receive Hep A.

Hepatitis B (Hep B)

Inactive virus: Transplant patients and household members may receive Hep B.

Inactive Polio Virus Vaccine (IPV)

Inactive virus: Transplant patients and household members may receive IPV instead of OPV (oral polio vaccine).

Live (attenuated) Vaccines: (should NOT be taken after transplant)

Varicella Zoster (Shingles)

Live virus: Transplant patients should not receive this vaccine. Family members may receive the vaccine.

Varicella (Chicken Pox)

If the family member breaks out with vesicles (pox), the transplant patient should not touch them. Rarely, a child who has received the vaccine may break out with a few vesicles. If this occurs, the pox should be covered with a band-aid and contact the transplant team for further guidance.

Measles, Mumps, Rubella (MMR)

Live Virus: Liver transplant patients should NEVER receive MMR after transplant. It may be given to siblings and other household members.

Oral Polio Virus Vaccine (OPV)

Live virus: Transplant patients and household contacts with patients should not receive OPV. Vaccine strains are transmissible. Because OPV is shed in the stool, it is recommended that IPV be given instead to transplant patients and those in frequent contact with them. Transplant patients should not change diapers of children given OPV!

Immunizations (Vaccines) for Pediatric Patients

Vaccines	Allow to use		Comments
	Transplant Recipient	Family Members	
Diphtheria/Tetanus toxoid/pertussis vaccine (DPT)	Yes	Yes	
Human Papillomavirus (HPV)	Yes	Yes	
Hemophilus B (HIB)	Yes	Yes	
Hepatitis A (Hep A)	Yes	Yes	
Hepatitis (Hep B)	Yes	Yes	
Oral Polio Virus Vaccine	No	No	Live vaccine strains can be transmissible as OPV is shed in the stool, recommend IPV for patient and their family member
Inactive Polio Virus Vaccine (IPV)	Yes	Yes	
Measles, Mumps, Rubella (MMR)	No	Yes	Live vaccine
Flu Shots	Yes	Yes	Never receive the nasal flu mist – this is a live vaccine
Pneumococcal (PCV-13 or PPVS-23)	Yes	Yes	
Meningococcal (Menactra)	Yes	Yes	
Tetanus	Yes	Yes	
Tuberculosis (TB) testing using purified protein derivative (PPD)	Yes	Yes	
Varicella (Chicken Pox)	No	Yes	Live vaccine. Transplant patient should not touch breakout vesicles (pox) of children or family member
Varicella Zoster (Shingles)	No	Yes	Live vaccine

Nutrition Following Liver Transplantation

Nutrition plays a key role in your recovery after liver transplantation. As with any surgery, adequate calories, protein, vitamins, and minerals are needed for wound healing. Also, your nutrient and diet needs may change if you have complications and/or side effects from your medications after your transplant.

Once you have successfully recovered from the early stage after transplant (for example: your appetite is back to normal, your weight is stable, and your wounds are healing well), then you should change to a diet that is low in saturated fat and high in fiber, fruits, and vegetables. This type of diet will help reduce risks for other chronic diseases including heart disease and diabetes. Always discuss your individual dietary needs and concerns with your dietitian or transplant provider (NP, PA, or doctor).

It is important to maintain a healthy weight. Obesity, gaining an unhealthy amount of excess fat, increases your risk of chronic diseases and can damage your new liver.

Short Term Nutrition Management

In the first month after your liver transplant, you will be recovering from the stress of surgery. It is important to take in enough protein, calories, vitamins and minerals to help your wounds heal and prevent muscle loss. Limit concentrated sweets or foods high in salt to reduce stress on your recovering body functions. If you have nausea, poor appetite or abnormal bowel functions, ask your health care team for suggestions. It is common for patients to feel full after eating a portion of the meal. If this happens, you should try eating smaller portions 6 times a day rather than larger portions 3 times per day.

Calories

You should eat enough calories to prevent muscle wasting and allow for gradual regaining of lean body weight that is often lost with severe illness before transplant and during hospitalization (bed-rest) after transplant. Choose nutrient-rich foods (dairy, whole grains, plant foods) instead of foods considered “empty calories” (for example: sodas, candy). If you have a poor appetite, broaden your scope of food choices to allow for food preferences. Fruits and vegetables need to be a part of your diet to provide enough vitamins and minerals.

Protein

Protein is important to promote healing and muscle gain. Extra protein is needed to overcome muscle breakdown caused by the high doses of prednisone given right after transplant .

Every day you should eat some foods that are highest in protein. Meats and fish have the highest concentrations of proteins, but many vegetarian sources are also excellent healthy choices that can add variety.

Animal-based products high in Protein	Plant-based products high in Protein
<ul style="list-style-type: none">• Fish• Poultry (chicken, turkey)• Egg whites (egg substitute)• Eggs (no more than 4 yolks per week)• Dairy products (milk, cheese, yogurt)• Red meat (beef, pork, veal)	<ul style="list-style-type: none">• Unsalted nuts• Peanut butter (or other nut butters)• Soy products (tofu, soy milk, tempeh)• Dried beans, lentils (split peas, kidney, black, pinto, navy beans, etc.)• Seitan (wheat gluten)

Choose lean protein foods more often and prepare them using lean cooking methods (for example, grill or bake meats rather than frying).

Sugar

Prednisone can decrease the uptake of sugar from your blood into your body's cells. This may cause your blood sugar (glucose) level to rise. This condition is called "hyperglycemia" (high blood sugar) or "steroid-induced diabetes". You can help manage this side effect by avoiding foods that are concentrated with carbohydrate (sugars).

Foods High in Simple Sugars and should be limited:	
<ul style="list-style-type: none">• Sugar• Molasses• Doughnuts, pastries, sweet rolls• Pies, cakes, cookies• Candy, chocolate candy• Ice cream, frozen yogurt, sherbet	<ul style="list-style-type: none">• Honey• Syrups (corn, maple, etc.)• Jam, jelly, marmalade (small amounts OK)• Soft drinks (diet sodas OK)• Other sweetened beverages (fruit juices, teas: diet OK)• Jell-O

If you develop high blood sugar, your diet may need to include the following guidelines:

- Eat at least three meals each day at regular, evenly spaced times. Avoid skipping meals. Each meal and snack should include some high quality protein. For example, a turkey sandwich on whole grain bread with salad. A snack could be 1/2 cup of 1% cottage cheese with fresh fruit or berries.
- Carbohydrate foods should be high in fiber. Choose to eat more whole grain products, legumes, and vegetables instead of refined white breads and cereals.
- Limit fruit to one serving per meal. One serving = 1/2 cup of chopped or canned fruit, or one medium-sized piece of fruit.
- Read food labels. Avoid foods with added sugars listed in the ingredients. Different names for sugars include: fructose, maltose, dextrose, corn syrup, high fructose corn syrup, honey, raw sugar, brown sugar, etc.

There are many resources available to learn how to best manage blood sugar. Ask your dietitian or healthcare provider for reading and education materials. Many healthcare providers or insurance plans provide diabetes management classes. Ask if they may be appropriate for you.

Sodium (salt)

Surgery and medications such as prednisone can cause your body to “hold on to” (retain) sodium and water. This can then cause increased blood pressure. To help avoid this, it is best to restrict salt intake. The “No Added Salt” diet, which is usually recommended, suggests eating less than 3000 mg of sodium each day. To follow these recommendations, you should:

1. Limit salt when cooking. Use herbs and spices for flavor.
2. Do not add salt after food is prepared.
3. **Avoid or limit the following foods:**

High salt soups

Most canned, dehydrated, and bouillon soups are high in salt. Read labels to choose those lowest in salts. Many popular brands now carry low sodium versions of soups.

Processed meats

Ham, bacon, sausage, frankfurters, cold cuts (bologna, salami, pastrami), corned beef, smoked or dried meat or fish, canned meat (unless labeled “no salt added”), salt pork.

Processed / canned food products

Cheese, packaged meals /dinner, most canned foods (except those labeled “no salt added”), frozen vegetables with sauces, pickles, sauerkraut, canned vegetable juice.

Restaurant items

Average amount of sodium per entree is 3000 mg.

Salted snacks

Salted crackers, pretzels, potato chips, corn chips, tortilla chips, popcorn, nuts, seeds, etc.

Condiments and miscellaneous:

Use in moderation: catsup, prepared mustards, horseradish, sauces (barbecue, chili, steak, Worcestershire, etc.), MSG (monosodium glutamate), soy sauce, meat tenderizers, pickle relish, olives, garlic salt, onion salt, salad dressings .

Note: salt substitutes are high in potassium and should only be used with your healthcare provider's permission.

Potassium

Some drugs such as Cyclosporine or Tacrolimus can increase your blood potassium levels. Other drugs such as Furosemide (Lasix®) can decrease blood potassium. Abnormal blood potassium levels can cause problems with muscle and heart function. Your health care team may change your medications and/or diet to help manage your blood potassium levels.

Below is a list of common foods high in potassium. You may be asked to either increase or decrease your intake of these foods. Remember that these are general guidelines and do not include all foods that contain potassium. If your diet needs to be stricter, ask your dietitian for a more complete list.

Fruits	Vegetables	Juices	Other Foods
<ul style="list-style-type: none">• Apricots• Avocados• Bananas• Dried fruit• Melons• Nectarines• Oranges• Peaches	<ul style="list-style-type: none">• Leafy greens• Pumpkins• Potatoes• Split peas• Dried beans• Lentils• Tomatoes	<ul style="list-style-type: none">• Orange juice• Prune juice• Tomato juice• V-8 juice	<ul style="list-style-type: none">• Milk and dairy• Peanut Butter• Nuts• Chocolate

Nutrition for Pediatric Patients Following Liver Transplant

Children and adolescents who receive liver transplantation may have had weight loss or delayed weight gain and growth before surgery. You probably had to make adjustments in your child’s diet to reduce the complications or discomfort of chronic liver disease. After surgery, however, the emphasis is to provide enough calories, protein, and other essential nutrients to promote healing, weight gain and growth. Encourage your child to eat a variety of foods and try to limit the amount of dietary restrictions you put on your child. However, your child’s diet should be somewhat low in sodium (salt) and concentrated sweets to reduce the body’s stress from medications and recovery from surgery.

The dietitian will work with you and your child to identify wholesome and tasty foods that are easily tolerated and good substitutes for fast food. Some infants and children may need tube feedings after transplant for normal growth and development. If nasogastric tube feeds are necessary, your healthcare provider will prescribe them and you will receive instructions on how to manage the feedings at home.

Long Term Nutritional Management

Transplant patients have increased risk for excessive weight gain, heart disease, diabetes, and low bone mineral density (osteopenia or osteoporosis) in the months and years after surgery. Some medications can be involved in the development of these complications, but researchers and healthcare providers believe diet and activity behaviors are very important. Risks can be reduced by adopting a low fat, low saturated fat, low cholesterol, high fiber “Heart Healthy” diet as outlined:

Foods	Choose more often	Choose less often
Vegetables	Minimum 3 servings each day. Try fresh or frozen that are prepared plain or with small amounts of unsaturated fats.	Vegetables prepared with butter, cheese or cream sauce.
Fruits	Minimum 2 servings each day. Fresh, frozen, canned, or dried.	Fruit juice; fruit served with butter or cream sauce.
Breads, Cereals, Grains, Dry Peas, Beans, and Soy	Whole grain bread, cereals, pasta and rice; dried peas, beans; baked goods made with unsaturated oil or margarine; soy foods.	Refined, white breads, cereals and baked good; white rice; dried peas or beans made with cream, butter, or cheese sauce.
Eggs and Dairy Products	Egg whites and substitutes; skim or 1% milk, yogurt and cottage cheese.	Egg yolks; whole or 2% milk, yogurt and cheese.
Meat and Poultry	Lean cuts with fat trimmed; turkey or chicken without skin; lean luncheon meat; cooking methods to help reduce fat.	Fatty cuts of meat; bacon and sausage; organ meats; any fried meat; high fat luncheon meat (salami)
Fish and Shellfish	All, cooked using low-fat methods.	Fish and shellfish cooked with butter, cheese or cream sauces.

Foods	Choose more often	Choose less often
Fats and Oils	Limit all fats to <30% of total calories. Margarine (made from unsaturated oils), including reduced- or non- fat varieties; reduced or non-fat salad dressings (including mayonnaise); liquid cooking oils; unsalted nuts and seeds.	Tropical oils (palm, palm-kernel, coconut); transfatty acids (hydrogenated fats); butter, lard, bacon fat, shortening, full fat salad dressings; coconut.
Sweets and Snacks <i>*Remember that while some of these products are low in fat, they still contain calories, mostly from sugars.</i>	Low- or non- fat frozen desserts and baked goods; flavored gelatins; candy such as gum drops; low salt pretzels and baked snack chips; air-popped popcorn.	Candy and baked goods made with butter, cream or tropical oils; high fat ice cream and frozen desserts; doughnuts; regular snack chips.

An easy way to think of this is to eat lots of vegetable/plant foods (whole grains, fruit, vegetables) and fish. Limit the amount of high fat animal foods. Read food labels carefully to avoid unnecessary fats and sugars.

It is also important to get enough calcium in your diet. Adults after transplant should have approximately 1200 mg of calcium (but less than 2000 mg) each day. This amount may be difficult to achieve if you don't eat dairy foods. Talk with your healthcare provider or dietitian about adding calcium supplements if you feel your dietary calcium is low.

Low-Fat Cooking Tips

- Microwave, broil, grill, or steam without adding fat.
- Use nonstick sprays or cookware.
- Trim all visible fat off meat and remove skin from poultry. (It is OK to remove skin after cooking)
- Experiment with reducing the amount of oil you use in baking. (Often recipes will taste just as good with less oil.)

There are many books and magazines devoted to health and nutrition. Enjoy trying new low fat, healthy recipes for the whole family.

Food Safety After Transplant

You are on medications that lower your immune system to prevent rejection of your transplanted liver. These medications increase your risk for foodborne illness. The following food safety tips can help when preparing, handling, and consuming foods.

- Wash hands and surfaces often.
- Avoid cross-contamination by washing cutting boards, utensils, dishes, and countertops often while preparing food.
- Wash all produce under running water, including those with inedible skins.
- Refrigerate foods promptly. Toss any foods that smell or appear spoiled.
- Cook to safe temperatures. Foods are safely cooked when they are heated to the USDA-FDA recommended safe minimum internal temperatures as noted below:

USDA-FDA recommended safe minimum internal temperatures

USDA-FDA recommended safe minimum internal temperatures				
Steaks/Roasts, Pork or Lamb Chops	Fish	Ground Beef, Pork, Lamb	Egg Dishes	Turkey, Chicken & Duck (whole or ground)
145°F	145°F	160°F	160°F	165°F

Food	High risk foods to AVOID	Lower risk foods to choose
Seafood	Raw or undercooked fish or shellfish, (for example, oysters)	Well-cooked seafood or fish (should be firm and flaky)
Red meat and poultry	Raw or undercooked red meat and poultry	Well-cooked red meat or poultry
Milk	Unpasteurized dairy	Pasteurized milk
Eggs	Raw or undercooked eggs or food containing them	Well-cooked eggs (firm yolks and whites), pasteurized eggs/egg products
Cheese	Soft cheeses made from unpasteurized milk (for example: moldy cheeses such as brie or camembert)	Hard cheeses, processed cheese, cream cheese, mozzarella, or soft cheese labeled “made from pasteurized milk”
Hot dogs and deli meats	Hot dogs and deli meats that have not been reheated	Hot dogs and deli meats reheated to steaming hot
Vegetables	Unwashed fresh vegetables	Washed fresh vegetables and cooked vegetables

See the USDA-FDA website (fda.gov) for further information on food safety for transplant recipients.

Fitness For Life

At UCSF, we strongly feel that regular physical activity is necessary after transplant for you to become as healthy and physically fit as possible. Everything you do to improve your overall health increases your chances of having a long life with your transplant. With the increasing shortage of organs available for transplant, it is a good idea to make healthy lifestyle choices . . . a second transplant may not come easily .

There is enough scientific evidence to say that regular physical activity:

- Decreases risk of death from heart disease.
- Prevents or delays the development of high blood pressure.
- Reduces blood pressure in people who already have high blood pressure.
- Keeps muscles and joints strong and functioning.
- Helps bones develop during childhood and helps adults prevent osteoporosis (thinning of the bones).

Special Importance of Physical Activity for Transplant Recipients

Long periods of inactivity and/or bed rest causes loss of muscle mass and strength (including strength of the heart) and limits your ability to perform exercise. A successful transplant does not automatically return you to normal physical activity. For muscles to regain their function and strength, they must be used regularly. Many people after transplant have high blood pressure, high cholesterol, and gain fat weight. All of these increase your risk of heart attacks or stroke.

You may resume physical activity 6 weeks post liver transplant. You can discuss with your transplant NP/PA or your local MD.

The weight gain following transplant is not due to prednisone. The weight gain is a result of the balance between how much and what types of foods are eaten and how many calories are burned through physical activity and your resting metabolism, which decreases with age.

How to Start an Exercise Program

When you decide to start your exercise or physical activity program, you should consider several things:

- type of exercise
- how often you will exercise (number of times per week)
- how long you will exercise each session (minutes per session)
- how you will continue to develop your program
- how hard you need to work or exert yourself

The key is to start slowly and increase gradually. That way your body will adapt to the exercise so you will feel comfortable and not get discouraged. You will definitely see progress as long as you stick with a program.

TYPE of Activity:

The best activities use large muscle groups in a rhythmic manner: walking, aerobics, cycling, swimming, jogging, machines such as climbers, stationary cycles, and ski machines. Choose an activity that is convenient and fun for you. There are no activities that are restricted for transplant patients.

If you have problems with your joints, an activity that supports your body weight and is not jarring will be best for you (such as cycling). If, after starting you feel discomfort in your joints, you may want to change your activity to one that is 'easier' on the joints such as swimming. Walking is easy, requires no special equipment other than comfortable, supportive shoes, and may be done anywhere, anytime.

FREQUENCY of Activity:

You should try to schedule your activity so you are doing something at least 5 days per week. Planning activity into your daily schedule will help you stay with it on regular basis. There is no time of day that is better for exercise . . . it all depends on what works for you and when you will be able to do it regularly.

DURATION of Activity:

Start with your activity for 10 minutes and gradually increase by 5 minutes every week to a goal of 30 or more minutes per session. **If** you are working to reduce or control your weight you will need to burn more calories, which means that you must exercise for a longer duration each session . . . the longer you exercise, the more calories you will burn.

INTENSITY of Activity:

The question is "how hard must I work myself when I exercise?" You should work hard enough to:

- make your heart beat faster
- allow you to breathe in a way that you can talk with someone during your session
- make you feel warm (perspiration is ok)
- feel that the exercise is "somewhat hard" for you

The increased effort can be worked in for short times during your exercise session. For example: climb hills and then walk down, or walk much faster for a block or two then walking slower for a block or two, etc. This will add variety to your session as well as further improve your fitness.

PROGRESSION of Activity:

It is best to start slowly and increase gradually. Start with a short duration of activity and gradually increase the time of each session. Start with a comfortable, lower level of intensity (work) and gradually increase the exertion level as you get used to the exercise. People may have problems, such as injuries, when they do too much exercise too soon. Gradual progression will prevent those kinds of problems.

Resistance (Weight) Training

Exercises that can easily be done at home to strengthen muscles include:

- climbing stairs
- standing up and sitting down into a chair (repeat 10–15 times, using your hands as little as possible so your legs do the work)
- lifting soup cans
- push-ups against the wall or from a counter

As you become stronger, you may want to use free weights or weight machines at a gym. Start with a low weight that you can comfortably lift 10–12 times. Remember that the weight being lifted will increase the stress on the joint (such as elbow or shoulder), so if the weight is too heavy or the exercise is not performed correctly you may hurt yourself. Don't hold your breath when lifting weights.

The CHALLENGE = Sticking with it!!

1. You must work hard enough with your body (heart, lungs, muscles, etc.) so that changes can happen.
2. You must stay with the program.

The fact is: increased physical activity is a personal commitment. No one else can do it for you and your health over time is dependent on being physically active on a regular basis.

U.S. Transplant Games

The U.S. Transplant Games is a wonderful celebration open to all recipients of solid organ and bone marrow transplants in the United States. All transplant recipients are welcome, whether or not they are athletes. The Games provide competition for those interested in challenging themselves and is also a chance to meet others from all over the country who have the similar experience of transplantation. At the first event in 1990, there were 400 recipients, and it has grown to over 5,000 recipients, family and friends. It is an exciting, inspirational event for everyone. The UCSF Transplant Service encourages anyone who is interested in participating.

The National Kidney Foundation (NKF) organizes the U.S. Transplant Games every 2 years. All transplant recipients (heart, lung, liver, kidney, pancreas, and bone marrow) of all ages and all athletic abilities are welcome and encouraged to participate. This is your time to show your family, friends, donors, and donor families that you're not "just" alive, but living life, too! Events include both team and individual races, swimming, golf, tennis, table tennis, bicycling, bowling and more.

There are also social events: opening and closing ceremonies, parties, workshops for donors, and a special tribute to all organ and tissue donors. For more information about the Transplant Games, please visit the National Kidney Foundation (www.kidney.org) or call the Northern California Chapter of the Kidney Foundation, 800-622-9010. Contact the American Liver Foundation for liver walks (www.liverfoundation.org) 800-GO LIVER.

Health Maintenance

Skin Cancer

Transplant patients have an added risk of cancer, especially skin and lip cancers. This risk increases with time due to long term immunosuppression (anti-rejection medications).

Immunosuppressive drugs make your skin sensitive to the sun causing you to burn easier, faster, and to a much greater degree than before your transplant. Prolonged, repeated exposure to the sun produces permanent skin changes and damage. Remember that you can burn even if your skin is dark because your medications may make you more sensitive to the sun.

Your risk of developing skin cancers increases with time after transplant. These cancers are ten times more common in transplant patients than in the general population. Therefore it is essential that you limit your exposure to the sun and that you protect yourself when you are outside.

Warning signs of skin cancers are any sores that bleed, scab, grow, or do not go away in a few weeks. Such sores are most likely to appear on the exposed parts of your body—face, neck, head (especially if you are bald), and your hands and arms. A mole that bleeds or changes color or size must be examined immediately. You must contact your local provider to be evaluated by a dermatologist. You should see a dermatologist yearly or more frequently if indicated. You may need to be referred to a “high risk” dermatologist. Contact your Transplant provider for more information.

The rays of the sun that are responsible for skin cancers are ultraviolet rays (UV rays) which are present even on cloudy days and in shady areas. We advise you to always protect your skin from exposure to these rays. Avoid the mid-day sun (10am–3pm) when ultraviolet rays are strongest. For extra protection from the sun wear sunscreen with SPF of least 30, lip balm, broad brimmed hats, long sleeves, and slacks every time when you go outdoors.

Some studies have shown that Vitamin B3 can decrease the incidence of squamous cell carcinoma. If you want, you can take it. It is ok to take Vitamin B complex daily.

Routine Self Examination

Cancers are more common when people are on immunosuppressive medications. Because of this, we recommend a monthly self-examination (ask your primary care provider if you need instruction how to do breast or testicular self-examination) and routine medical checkups. PAP smears and breast exams should be done every year. Reports should be sent to the Transplant Office. Your local physician can perform these tests.

Acne

Prednisone may cause acne on your face, chest, shoulders, or back. Cyclosporine can make your skin oily in these areas.

If your skin becomes excessively dry, temporarily stop using soaps on these areas to allow the skin to recover its natural moisture. You may use over the counter preparations (such as Oxy-5 or Oxy-10) to treat acne. We recommend that you do not use Retin-A (a powerful acid form of Vitamin A). It causes increased sun sensitivity which should be avoided.

If acne remains a problem, tell us about it. Severe or infected acne must be treated by a dermatologist.

Dry Skin

If you have problems with dry skin, use a mild soap and apply body lotion after bathing. For severe dry, scaly skin you may need a good body lotion. There are many good products available from your local drug store without a prescription.

Cuts and Scratches

Keep minor cuts and scratches clean and dry by washing daily with soap and water. If you wish, you may apply an antiseptic such as Betadine solution. For large cuts, dog bites, etc., contact your doctor immediately. You will not need any special skin care unless you have problems with acne or dry skin. You should take a bath or shower as often as necessary (daily or every other day) to keep your skin clean. Use any soap that you prefer.

IMPORTANT: Any persistent skin problem requires an evaluation by a dermatologist.

Increased Hair Growth

This problem is caused by Cyclosporine and is especially annoying to women. You can remove hair with a hair removal cream. There are several creams on the market made especially for your face. Be sure to follow bottle directions concerning testing. Hair removal cream causes severe irritation to the eyes, mucous membranes, and lips so apply carefully. A safer way to diminish the look of excessive hair is to bleach the extra hair growth with 50% peroxide solution. When your Prednisone dose is less than 20 mg per day, you may want to consider waxing or electrolysis for permanent hair removal.

Caution: Parents should be particularly careful when using these products on infants and young children since their skin tends to be more sensitive.

Hair Loss

Tacrolimus® (Prograf) can cause hair loss. This usually stops or slows down as you have your dose lowered.

Hair Care

Prednisone often affects the condition of your hair because it makes your hair weaker. Permanent wave lotion, tints, dyes, and bleaching may cause your hair to break. We recommend that you wait until your Prednisone dose is less than 20 mg per day to have a perm or to color your hair.

Dental Care

Routine dental work should be done six months after your transplant. This includes having your teeth cleaned. However, if you have tooth or gum pain you should see your dentist immediately.

- You will not need to take an antibiotic when you have dental work done, unless your dentist deems necessary due to a possible infection.

We follow the American Heart Association (AHA) guidelines for prophylaxis. Your local healthcare provider and dentist are aware of these guidelines and should refer to them should they have questions regarding antibiotics for dental work in transplant patients.

It is extremely important that you practice good mouth care after your transplant to maintain healthy gums and teeth, as well as to avoid infection. Brush your teeth once or twice a day. Floss your teeth once a day; be gentle if you notice any bleeding. Contact your dentist or dental hygienist if your gums continue to bleed with flossing.

If your dental work was not completed before transplantation, you must see your dentist for evaluation within the first few weeks after you return home.

Follow the procedure outlined below when you visit the dentist:

- Make an appointment.
- Inform the dental office that you have had your transplant.

The Transplant Center will fax or mail a dental clearance letter as requested by the dentist or by you. See sample dental letter on the next page.

Prednisone Dose and Routine Dental Care

Your Prednisone dose should be 5 mg once/day (or less) when dental work is done to reduce the possibility of infection and bleeding, and to increase the healing process. But, do not avoid dental care if your Prednisone dose is high and you have tooth or gum pain. It is better to have healthy teeth and gums than to allow them to become infected.

If your dentist has any questions or if you have any questions, contact the Liver Transplant Office for further information.



University of California
San Francisco

Liver Transplant Program

UCSF Box 0780
505 Parnassus Ave., Room M-896
San Francisco, CA 94143

tel: 415.353.1888
fax: 415.353.8709

www.ucsf.edu

Date _____

Name _____

DOB _____

To Whom It May Concern:

Please be advised that this patient received a liver transplant on _____.

There are no contraindications for dental work at this time. Please see the following guidelines for post liver transplant patients.

Antibiotic prophylaxis: We follow the American Heart Association guidelines for antibiotic prophylaxis . Post liver transplant patients do not need prophylactic antibiotics for routine dental procedures. Antibiotics may be given at the discretion of the treating Dentist if required as standard of care for a particular procedure .

Local Anesthetic: This patient may receive local anesthesia such as Novocain as needed.

Pain Medications: The patient should not receive products containing aspirin or ibuprofen. The patient may receive products containing Tylenol as long as the total daily dose does not exceed 2 grams per day. Patients may receive a short course of narcotics if needed.

We can be reached at (415) 353-1888 for any additional questions.

Thank you,

UCSF Transplant Service

Smoking

Don't smoke. You stopped smoking (or were supposed to) prior to transplant. Now there are even more reasons not to smoke.

Smoking damages the lungs, putting you at great risk for lung infections including bronchitis, emphysema, and pneumonia. It also increases your risk of head & neck and lung cancer **(especially after transplant)**.

Smoking lessens the ability of your red blood cells to carry oxygen, so less oxygen goes to all of your tissues decreasing your ability to heal.

Smoking narrows your blood vessels, especially those in your legs, arms, and heart.

Smoking also increases acid in your stomach that can delay or prevent the healing of any ulcers that may occur.

All of these problems are more serious in someone taking immunosuppressive medications.

Talk with your primary care provider about short term medications or smoking cessation programs that may be available if you need help to stop smoking.

Alcohol and Illegal drugs

Do not drink alcoholic beverages or take drugs, including marijuana. This includes non-alcoholic beers and wines. (They still contain small amounts of alcohol). Alcohol is metabolized (broken down) by the liver and causes liver damage as well as liver failure. Almost all medications are metabolized by the liver. The combination of these medications with alcohol could seriously harm your liver.

Smoking marijuana exposes your lungs to harmful fungi that can lead to life threatening infections.

If you are having trouble living without alcohol or drugs, you need to know that this can be a life threatening problem. Help is available. Contact your local primary care physician, your transplant provider, or the transplant social worker for help.

Sexuality

You may resume sexual activity when you feel like it. Remember to always practice safe sex if you are not in a monogamous relationship. Contact the transplant team to review immunosuppressive medications **before becoming pregnant or fathering children**. Contact your gynecologist to discuss methods of birth control. Some of your immunosuppression may cause birth defects; therefore, your medications may need adjustment before you become pregnant or before you father a child.

Family Planning

Some people want to have children after they have had a liver transplant. For many women this is possible, but there are also risks to the mother and fetus. It is important to discuss these risks with the Transplant

Team so that you can make the right decision for you and your family. Male transplant recipients should also re-view medications with the transplant team before fathering children. Some immunosuppressive medications may need to be changed or discontinued before pregnancy to protect the baby from potential birth defects.

Return to work/school/disability: frequently asked questions

1. When can I return to work / school?

Most patients are able to return to work 3–6 months after transplant with no restrictions. Generally, there are no weight lifting or activity limits approximately 3 months after transplant. Most people start planning about a month after transplant for their return to work. Of course, this depends on your particular recovery.

Children may be ready to return to school approximately 2-3 months after transplant. We will write a letter to your child's school that will include the following, as well as any special needs for your child:

- No P.E. for at least 3 months after transplant. Contact sports like American football, karate and wrestling, are prohibited
- Recess is encouraged
- Need for frequent lab work and clinic appointments

2. How long will I be on disability?

Disability benefits depend on your physical condition. If you were on disability at the time of your transplant, your continued eligibility for benefits will be reviewed some time after your surgery. The timing of this review will depend on your particular disability plan. If your liver function is good and your condition is stable, the liver transplant team will not be able to say that you continue to be disabled. Generally, by 6 months after transplant, patients are no longer disabled because of transplantation.

3. What should I do with disability forms?

The transplant team will be able to complete your disability forms within the first 3 months after transplant. You may bring the forms to your clinic appointments or mail them to your transplant provider. Generally, after 3 months, your primary care provider will be seeing you more often than the transplant team and is the appropriate person to complete any disability forms.

4. What if I don't have a job?

If you have been off from work prior to transplant, there is help available to assist you in finding a job. The Department of Vocational Rehabilitation can help you identify your skills and options. They can also help you with job finding skills, re-training or schooling. Call the transplant social worker for more information.

Call **1-800-LIVER TX** (548-3789) with non urgent questions, or messages for the transplant team. Messages may be left 24 hours a day. Your message will be returned within one business day.

Travel

There are no travel restrictions after transplant if your condition is stable. See page 33 for additional information about immunizations. Contact your local health care provider or travel clinic for questions. When traveling, remember to plan to take enough medication plus extra so you will be prepared if you have unexpected changes in plans. Always carry medication on board rather than checking with your luggage.

More information can be found at www.cdc.gov. Remember that you may **NOT** receive live vaccines. (Please review pages 34 and 35 of this book.)

Other Health Care Concerns

Illness and Infection

You need to protect yourself from infection because your immune system is suppressed. Wash your hands with soap and water often to wash away bacteria and viruses. Keep your hands away from your face and mouth. Wash hands after using the bathroom and before eating. Avoid sick people. Ask sick friends to visit when they are well. If you have a wound and do your own dressing changes, wash your hands before and after you change the dressing. When you cough or sneeze, use tissues, discard them in the trash, then wash your hands.

Even when taking all possible precautions, there are times when you may get an infection anyway. If you have a fever over 100.4° F (38° C) or you do not feel well, see your local healthcare provider and call the transplant office. You may need to have blood drawn and a chest x-ray. You will be prescribed an antibiotic if you have a bacterial infection. Sometimes you may need to be admitted to the hospital for treatment.

Remember that stomach pains, problems with your heart, problems with breathing, or fevers cannot be diagnosed over the phone. See your own healthcare provider for any treatment. You may call us at 415-353-1888 for urgent questions about your care. Providers for pediatric patients should call 415-476-5892.

Call **1-800-LIVER TX** (548-3789) with non urgent questions, or messages for the transplant team. Messages may be left 24 hours a day. Your message will be returned within one business day.

Colds and Flu

You will get colds and flu. If you get a cold, you may take over the counter medications (ex: Sudafed, Dimetapp, etc.). Avoid cold remedies that contain aspirin and Ibuprofen. Limit Tylenol (acetaminophen) to 2000 mg per day. Read labels and prescriptions—Tylenol is contained in many medications. If you have a cough, you may take cough medicine (ex: Robitussin). However, if your cough becomes productive (with green phlegm or sputum) and you run a fever, call your local healthcare provider. You may need a chest xray and antibiotics.

For Children: Please check with your pediatrician.

Inform the Transplant Office whenever you are exposed to any communicable disease that you have not previously had. This is particularly important for liver transplant children.

Adults: 1-800-LIVER TX

Monday–Friday, 8am–5pm

Children: 415-476-5892

Herpes Zoster (Shingles)

Shingles appear as a rash or small water blisters usually on the chest, back or hip. They may also occur on other areas. The rash may or may not be painful. You should notify your primary care provider and transplant provider if this rash appears. (For vaccine questions, see page 34)

Persistent Headaches

Tacrolimus and Cyclosporine can cause headaches. Please inform your transplant provider for adjustments. If they become persistent you may need to see a neurologist.

Skin Eruptions

If you develop warts, persistent sores, or moles you must see a physician, preferably a dermatologist. If a skin eruption requires medical treatment, have your health care provider call the transplant office 1-800-LIVER TX to discuss the treatment. Pediatric patients should call 415-476-5892.

Thrush

This is an infection caused by fungus that usually affects the mouth. It can also affect the esophagus (the tube from your mouth to your stomach), the vagina in women or the diaper area in infants. With oral thrush, your mouth may be very tender and you may notice that your tongue is covered by a white film. Mycelex Troche or Nystatin (a yellow liquid) can help clear up this infection. It is helpful to eat soft foods or drink liquids. With vaginal thrush, you will notice an increase in vaginal discharge and possibly a change in its color. Your genital area may also itch. Vaginal creams and other medicines are available. Talk with your primary care provider or the Transplant Team if you think you have thrush. Do not take oral antifungals (fluconazole) without checking with your transplant team first. Antifungals can cause a drastic elevation in your tacrolimus or cyclosporine blood levels.

Chicken Pox, Measles, and Mumps:

Chicken pox is a virus that causes a mild fever and small red bumps on the skin and can be very itchy. It is spread by coming in contact with another person infected with the virus. Being in the same room with an infected person can transmit the virus. If you had chicken pox as a child, you most likely will not get it again. However, if you have not had chicken pox or you are not sure, let the Transplant Team know. This is a relatively mild disease in children. In adults chicken pox is more severe—especially in immuno compromised adults. If you suspect you have been exposed to chicken pox, call the Transplant Team immediately.

Emergency Medical Identification

You should wear a medical identification bracelet or necklace at all times after your transplant. You may obtain a form to order one at your local pharmacy. Indicate on the form that you have had a liver transplant, and are “immunosuppressed.” Include the 24 hour hotline number 415-353-1888 (for pediatric patients: 415-476-5892) to be printed on the tag.

It is a good idea to also have your local health care provider's phone number printed on the tag. If the transplant patient is a child, you may want to print the parents' first names and home phone number(s) as well.

Guidelines for Visiting Transplant Children

- We support and encourage parents or guardians to stay with their child. Most patient rooms have sleeping accommodations for one family member to stay overnight. Our family resource coordinator can assist with local housing for longer stays.
- Siblings of all ages are encouraged to visit.
- All parents, family members, and visitors must check in at the hospital information desk in the lobby. A brief security and health screening will be conducted.
- Visitors with any signs or symptoms of infection or illness, such as fever, cold, cough, sore throat, runny nose, diarrhea, vomiting, or any other cold or flu symptoms, are asked to postpone their visit.
- All children must wash their hands thoroughly before entering the patient's room.
- No children visitors may receive treatments or medications of any kind while they are visiting.
- As there is no care available for visiting children, please make arrangements for their supervision. The staff may ask visiting children to leave if they are unsupervised or disruptive.
- There are no accommodations for visiting children to spend the night.
- If there are any falls, bumps, cuts or accidents, a parent must notify the nurse promptly. You may be asked to take the visitor to the emergency room for treatment.
- Visiting children must leave the Pediatric floor immediately when the visit is over.

Thank you for being careful and following these guidelines. If you have any questions, please feel free to ask any of the nurses.

This page is intentionally left blank.

Resources

Getting Back Into Life!

A serious illness can create many personal and family stresses. After your transplant, there will be different types of adjustments. You and your family can expect some ups and downs related to your physical condition as well as emotional reactions to this new phase in your life. For example, you may feel somewhat anxious when you get your labs drawn or when you need to have tests if your lab values change. You or your caregivers may have mixed feelings as you become more independent and responsible for your own care. These are normal reactions.

If you have been sick for a long time, you may have had to give up some of your normal life such as child care, work, and activities with friends. You and people close to you may have some ups and downs as you get back to your normal activities and roles. Your illness may also have changed your relationship with your significant other. Sometimes it may help to talk to someone else who knows what you are going through. In most communities, there are transplant support groups. The transplant social worker is available to talk with you about these transitions. The social worker also has lists of the community support groups and other resources which may help you.

If you had a sudden, unexpected transplant, you may experience some sadness or depression as you start to feel physically better. Remember, you did not have a chance to plan or prepare for the changes in your life and you will need to do that as you recover. Again, please speak with the transplant social worker or the transplant team if you are having difficulty.

No one can predict exactly what your recovery will be like. Complications, minor and serious, may occur any time. Remember that the transplant team will work with your primary health care providers and you to take care of you.

There may also be some practical problems after transplant such as insurance or financial issues. Your medical care following transplantation is very expensive. Therefore, you must keep current with your insurance. Know what it covers, how it works, and what your responsibilities are. You must check with your insurance company to find out which laboratories and pharmacies you may use with your plan. Most insurance companies will only pay when you use the health care providers they approve. Remember, it is your responsibility to make sure any visits to UCSF have been approved (authorized) by your insurance. If your insurance coverage changes, please call **1-800-LIVER TX (548-3789)** to update your records.

Some people, as they recover after transplant, lose eligibility for disability based insurances like Medi-Care and Medi-Cal. If you think you are going to lose your insurance, talk with either the financial counselor or transplant social worker about what other coverage may be available.

Do not wait to call until you have no insurance. You must plan ahead.

Other practical issues that you should continue to keep updated include advance directives, wills, and power of attorney, especially if you have no legal next of kin.

There is light at the end of the tunnel!!

Life after transplant can be a real roller coaster ride for patients, families, and friends. The goal of transplantation is to help you return to good health and your usual lifestyle. Most patients are able to return to work, raise families, and lead full, physically active lives after transplantation.

Your transplant team looks forward to helping you in whatever way we can. Talk to us.

Adult Support Group

The Liver Transplant Support Group is held on Tuesdays after the adult Transplant Clinic on 9 Long in the Solarium from 1–2pm. It is open to all present and future liver transplant recipients and their families and friends.

Caregiver Support Group

The Caregiver Support Group is held on the first Friday of the month at 9 Long in the Solarium from 1-2pm. It is open to all caregivers only, no patients. If you would like more information, please call 415-353-1040.

Pamphlets and Organizations to Help You

Pamphlets available at www.liverfoundation.org:

- Biliary Atresia
- Decide to Make a Difference (organ donor)
- Diet and Your Liver
- Facts About Liver Transplantation
- Hemochromatosis
- Liver Disease: A Problem for the Child
- Viral Hepatitis: Everybody's Problems
- Your Liver Lets You Live

Organizations:

- **American Liver Foundation**
75 Maiden Lane
New York, New York 10038
1-800-GO-LIVER (465-4837)
www.liverfoundation.org

Northern California Chapter
One Sansome Street, Suite 2100
San Francisco, CA 94104
415-984-3169
- **Centers for Disease Control**
<http://www.cdc.gov>
(See travel guidelines)
- **Transplant Recipients International Organization (TRIO): Your voice of the transplant**
800-TRIO-386
info@trioweb.org
- **United Network for Organ Sharing (UNOS)**
www.unos.org
- **Donor Network West**
888-570-9400
www.donornetworkwest.org
(Pediatric)
- **Children's Liver Disease Foundation**
www.childliverdisease.org
- **Children's Organ Transplant Association**
800-366-2682
www.cota.org

Notes

Notes

Notes

Notes

Notes

